

Investigating the Relationship between Perceived Discrimination, Social Status, and Mental Health

Hedwig Lee¹ and Kristin Turney²

Abstract

A growing body of evidence suggests that experiences with discrimination have implications for mental health and that these associations may vary by social status. We use data from the Chicago Community Adult Health Study (CCAHS) to examine how two types of perceived discrimination, chronic everyday discrimination and major lifetime discrimination, are linked to mental health and how this association varies by race/ethnicity, gender, and socioeconomic status. Results indicate that everyday discrimination is generally independently linked to greater depressive symptoms, loneliness, and hostility across all social status groups. Major discrimination is not associated with depressive symptoms or loneliness after adjusting for a host of covariates but is associated with hostility, especially for certain groups. These findings highlight the need to examine multiple indicators of discrimination and mental health and to pay attention to both differences and similarities in these associations by social status.

Keywords

discrimination, mental health, race/ethnicity

INTRODUCTION

A growing body of research examines the consequences of perceived discrimination, a unique psychosocial stressor, for mental health (Paradies 2006; Williams and Mohammed 2009; Williams, Neighbors, and Jackson 2003).¹ Perceived discrimination is consistently, positively associated with impaired mental health across a vast array of cross-sectional and longitudinal data and has been linked to multiple mental health outcomes, including depressive symptoms, psychological distress, anxiety, and psychiatric disorders (Paradies 2006; Pascoe and Richman 2009; Williams and Mohammed 2009; Williams et al. 2003). However, little is known about variation in the association between perceived discrimination and mental health across social status groups such as race/ethnicity, gender, and socioeconomic status (SES), despite theoretical perspectives suggesting that both levels of stress and the association between stress and health

may vary across these groups in important ways (Pearlin et al. 2005; Thoits 2010; Turner and Avison 1989, 2003). Indeed, stress is unequally distributed across social status groups, with lower status groups such as blacks, women, and the economically disadvantaged facing disproportionately high levels of stress and having reduced resources to buffer against the negative consequences of stress (Pearlin et al. 2005; Sapolsky 2005; Thoits 2010; Turner and Avison 2003).

Guided by the stress theory perspective, which emphasizes inequalities in the distribution of vulnerability to stress across demographic groups, we

¹University of Washington, Seattle, WA, USA

²University of California Irvine, Irvine, CA, USA

Corresponding Author:

Hedwig Lee, Department of Sociology, University of Washington, 211 Savery Hall, Seattle, WA 98195, USA
Email: hedylee@uw.edu

examine variation in the mental health consequences of discrimination across social status groups. Using data from the Chicago Community Adult Health Study (CCAHS), a multiethnic population-based sample of adults, we answer two research questions. First, how is major acute perceived discrimination and chronic perceived discrimination independently associated with depressive symptoms, loneliness, and hostility? Second, how does the relationship between perceived discrimination and mental health vary by race/ethnicity, gender, and SES? Our diverse sample allows us to estimate variation in this relationship across demographic groups. Considering that mental health problems are associated with social and economic disadvantages throughout the life course—including financial hardship and difficulty in personal relationships—understanding the antecedents of impaired mental health is important (Coyne 1976; Miech and Shanahan 2000).

BACKGROUND

Perceived Discrimination as a Unique Stressor and Its Dimensions

Stress theory is a useful theoretical framework that helps explain both the unequal distribution of stress across the population and the differential association between stress and health across social status groups (Pearlin et al. 2005; Thoits 2010; Turner and Avison 2003). Stressful life events often arise from distinctive social contexts that characterize the lives of lower status groups such as racial/ethnic minorities, women, and the economically disadvantaged, and it follows that disadvantaged groups experience greater stress than their more advantaged counterparts. Lower status groups also lack the resources to cope with stressors (e.g., social support, leisure time, or financial resources to engage in stress-relieving activities) and thus may be more vulnerable to mental health impairments (Pearlin et al. 2005; Sapolsky 2005; Thoits 2010; Turner and Avison 2003). Stressors that are related to ascribed social statuses such as race/ethnicity, gender, or SES may be especially detrimental for mental health, as they often lead to additional burdens for disadvantaged groups by limiting opportunities and threatening identity (Pearlin et al. 2005).

Previous research indicates that perceived discrimination is a unique psychosocial stressor with important implications for physical and mental

health inequalities (Clark et al. 1999; Pascoe and Richmond 2009; Thoits 2010). In fact, perceived discrimination and the unfair treatment associated with it is one of the most important classes of psychosocial stressors (Wethington, Brown, and Kessler 1995). Further, the effect of perceived discrimination on mental health is similar in magnitude to the effect of more commonly studied stressors such as job loss, divorce, and death of a loved one (Kessler, Mickelson, and Williams 1999). Much evidence linking perceived discrimination and mental health examines racial/ethnic discrimination among samples of blacks and other nonwhite minority groups (Paradies 2006; Williams and Mohammed 2009). However, racial discrimination is only one of many forms of discrimination; individuals frequently experience discrimination based on other identifying status-based characteristics (Kessler et al. 1999). Both racial discrimination and discrimination based on other identifying characteristics can lead to mental health impairments (Kessler et al. 1999; Thoits 2010).

Similar to other stressful experiences, discrimination is a multidimensional construct, occurring in multiple domains (e.g., places of employment, restaurants, and neighborhoods) and over time. Examples of major acute discriminatory experiences include being unfairly fired from a job or stopped by the police, and chronic or “everyday” discriminatory experiences (sometimes referred to as “chronic daily hassles” or “day-to-day experiences”) are characterized as frequent irritations and indignities in everyday situations (e.g., being treated with less courtesy than others) (Williams et al. 2003). Although discrimination research most commonly assesses the consequences of major acute discrimination and chronic discrimination separately, there is evidence that these different dimensions of discrimination often co-occur and the relative consequences of them may vary (Bennett et al. 2010; Pascoe and Richman 2009; Williams et al. 2003). Individuals in disadvantaged groups are particularly likely to experience both acute and chronic stressors, and understanding how these stressors are independently associated with mental health can provide insight into the mechanisms linking discrimination to health (Williams et al. 2003). In addition, discrimination often co-occurs with and is compounded by other stressors such as familial conflict, financial strain, and stressful life events (e.g., exposure to violence and death of a spouse or child), leading to chronic

and repeated exposures to hardships and stressors across the life course, particularly among vulnerable populations (Kessler et al. 1999; Pearlin et al. 2005; Thoits 2010; Williams et al. 2003).

Previous Research on Discrimination and Mental Health

A large body of research has examined how perceived discrimination is associated with adverse mental health such as anxiety, depression, fear, frustration, helplessness, hopelessness, paranoia, resentment, and self-esteem (Burgos and Rivera 2009; Clark et al. 1999; Paradies 2006; Rivera et al. 2011; Williams and Mohammed 2009; Williams et al. 2003). Although the majority of this research has used mono-racial samples (often comprised of only blacks) and/or convenience samples that cannot be generalized to other groups, the association between perceived discrimination and poor mental health is also found in population-based multiethnic and multiracial samples (D'Anna, Ponce, and Siegel 2010; Kessler et al. 1999; Williams et al. 1997).

Though the association between discrimination and some mental health outcomes (e.g., depressive symptoms) is well documented, much less is known about how discrimination is associated with outcomes such as loneliness and hostility. Loneliness itself can be characterized as one possible indicator of the larger concept of social exclusion. Research on social exclusion makes clear that discrimination, stressful events, and exclusion from society, particularly due to disadvantaged social status, are bound up together in a life course of exposures that have an adverse impact on mental health (Brown, Bhrolchain, and Harris 1975; Brown and Harris 1978; Ross and Mirowsky 1989; Thoits 1982, 2010). Previous research suggests chronic social and marital stresses are associated with greater loneliness (Hawkley et al. 2008), and it is plausible that discrimination (another dimension of stress) operates similarly. Additional research documents how racial discrimination can lead to isolation of individuals in the workplace and in other social settings (Essed 1991; Feagin and McKinney 2003; Forman 2003). Other forms of systemic discrimination, such as residential segregation or political exclusion, could also lead to loneliness (Feagin and McKinney 2003; Williams and Mohammed 2009). However, to our knowledge, no previous

empirical research examines the relationship between perceived discrimination and loneliness.

Similarly, discrimination may lead to externalizing reactions such as anger and hostility. Anger can be an important and frequent psychological stress response resulting from race-based discrimination experienced by blacks (Brondolo et al. 2005; Clark et al. 1999; Feagin and Sikes 1994; Gibbons et al. 2010), as well as discrimination faced by other groups such as women (Swim et al. 2001). Indeed, perceptions of discrimination that provoke anger may lead to a coping response such as hostility or aggression (Clark et al. 1999; Feagin 1991). In addition, other research shows that anger and hostility (expression and suppression) may be an important pathway linking experiences with discrimination to substance abuse and/or increased blood pressure (Gibbons et al. 2010; Krieger 1990; Whitbeck et al. 2001). Research examining the relationship between racial discrimination and anger or hostility has been largely based on mono-racial samples (for exceptions, see Broudy et al. 2007; Rivera et al. 2011).

Variation in the Association between Perceived Discrimination and Mental Health

It is well known that certain population groups are more frequently exposed to discrimination. Racial/ethnic minorities, compared to whites, experience higher levels of discrimination (Borrell et al. 2010; Thoits 2010; Williams and Mohammed 2009). In addition, black men report higher levels of discrimination than black women (Borrell et al. 2010), which may result from them being stereotyped as threatening and thus experiencing more discriminatory incidents (Carter 2007). These gender differences have also been found among Hispanic populations (Borrell et al. 2010). With respect to SES variation in discriminatory experiences, current research comes to inconsistent conclusions (Brondolo et al. 2009; Dailey et al. 2010; Kessler et al. 1999).

Conceptually, a stressor such as discrimination may be more detrimental to disadvantaged groups such as racial/ethnic minorities, women, and those facing economic hardship. But relatively little research has examined the association between discrimination and mental health by social status. There are several exceptions. For example, Kessler and colleagues (1999) find that by and

large, the association between perceived discrimination and mental health is consistent across groups (also see McLaughlin, Hatzenbuehler, and Keyes 2010; Williams et al. 1997). Others, however, find that discrimination is more harmful to the mental health of Latina women than white women (D'Anna et al. 2010) or black Latinos than non-black Latinos (Burgos and Rivera 2009).

Even less research explores how the relationship between discrimination and mental health varies across SES (D'Anna et al. 2010; Forman 2003). On the one hand, discrimination might be more harmful to high-SES groups if these groups view discrimination as a threat to their social status. On the other hand, it might be more harmful for low-SES groups who may be treated more harshly due to their lower status and have fewer resources to cope with the stress of discrimination (Williams and Mohammed 2009).

Little research has examined the varying association between discrimination and mental health by race and gender (Williams and Mohammed 2009). Some evidence shows that despite higher reports of self-reported discrimination among minority men than women, the relationship between discrimination and depressive symptoms is stronger for black and Hispanic women compared to black and Hispanic men (e.g., Borrell et al. 2006; Flores et al. 2008). Indeed, discrimination may be particularly consequential to black and Hispanic women's psychological health due to their double minority status, burdened by experiences of both racial/ethnic and gender discrimination and occupying multiple roles within the labor force and the home (St. Jean and Feagin 1998; Troxel et al. 2003). Most of this evidence is drawn from mono-racial samples, so comparisons cannot be made across race and gender groups.

METHOD

Data Source

We use data from the Chicago Community Adult Health Study, a cross-sectional survey designed to examine the biological, social, and environmental correlates of adult physical and mental health. The CCAHS is a multistage probability sample of 3,105 adults (18 years and older) living in Chicago, stratified into 343 neighborhood clusters as defined by the Project on Human Development

in Chicago Neighborhoods (Sampson, Raudenbush, and Earls 1997). Face-to-face interviews were conducted with one respondent per household between May 2001 and March 2003, and the response rate was 72 percent.

Key Variables

Mental health. We examine three mental health indicators: depressive symptoms, loneliness, and hostility. Depressive symptoms is measured with a modified version of the Center for Epidemiologic Studies Depression Scale (CES-D), commonly used to measure distress (Radloff 1977). We average responses to respondents' reports of the following 11 symptoms in the past week (1 = *never*, 2 = *hardly ever*, 3 = *some of the time*, 4 = *most of the time*): I felt depressed; I felt that everything I did was an effort; my sleep was restless; I was happy; I felt lonely; people were unfriendly; I enjoyed life; I did not feel like eating; I felt sad; I felt that people disliked me; I could not get going ($\alpha = 0.85$). Loneliness, measured with a shortened version of the Revised UCLA Loneliness Scale, is the average of the following (1 = *never*, 2 = *rarely*, 3 = *sometimes*, 4 = *often*): I lack companionship; I feel left out; I feel isolated from others ($\alpha = 0.77$) (Hughes et al. 2004). Finally, hostility is an average of respondents' responses to the following (1 = *disagree strongly*, 2 = *disagree somewhat*, 3 = *agree somewhat*, 4 = *agree strongly*): Most people inwardly dislike putting themselves out to help other people; most people will use somewhat unfair means to gain profit or an advantage rather than lose it; no one cares much what happens to you; I think most people would lie in order to get ahead; I commonly wonder what hidden reasons another person may have for doing something nice for me ($\alpha = 0.74$) (Cook and Medley 1954). Higher values on each of the scales indicate worse mental health.

Perceived discrimination. We examine two indicators of perceived discrimination: everyday discrimination (Williams et al. 1997) and major discrimination (Kessler et al. 1999; Williams et al. 1997). The everyday discrimination scale assesses the occurrence and frequency of perceived chronic interpersonal discrimination that individuals experience. Respondents were asked to report how often the following occurs (0 = *never*, 1 = *less than once a year*, 2 = *a few times a year*,

3 = *a few times a month*, 4 = *at least once a week*): You are treated with less courtesy or respect than other people; you receive poorer service than other people at restaurants or stores; people act as if they think you are not smart; people act as if they are afraid of you; and you are threatened or harassed ($\alpha = 0.75$). We sum individuals' responses to the five questions. Major discrimination assesses the frequency of acute discriminatory events. Respondents reported if, at some point during their lives, they experienced the following (0 = *no*, 1 = *yes*): were unfairly fired from a job or were unfairly denied a promotion; were not hired for a job for unfair reasons; were unfairly stopped, searched, questioned, physically threatened, or abused by the police; and were unfairly prevented from moving into a neighborhood because the landlord or realtor refused to sell or rent you a house or apartment. As with everyday discrimination, we sum responses to the individual questions and higher values indicate more experiences of major lifetime discrimination.² The correlation between everyday and major discrimination is 0.42 ($p < .001$).

Key demographic variables. We adjust for demographic variables including race/ethnicity, gender, and household income. Race/ethnicity is measured with the following four dummy variables: non-Hispanic white (reference category), non-Hispanic black, Hispanic, and non-Hispanic other race (includes American Indian, Asian, and Pacific Islander). Respondents had the option to report more than one race; those who reported being white in addition to another race (e.g., black) were considered to be the other, nonwhite race (e.g., a respondent who reports being both black and white is considered black). A dummy variable indicates the respondent is female. Annual household income is measured with the following dummy variables: less than \$10,000; \$10,000 to less than \$30,000; \$30,000 to less than \$50,000; and \$50,000 or greater (reference category).

Additional control variables. Age, measured in years, is represented by the following dummy variables: 18 to 29 (reference category), 30 to 39, 40 to 49, 50 to 59, 60 to 69, and 70 and older. Immigrant status includes the following dummy variables: first generation, second generation, and third generation or higher (reference category). Education includes the following categories: less than 12 years, 12 years to 15 years, and 16 years or greater (reference category). A dummy variable indicates the respondent is currently working, and homeownership and living

alone are also dummy variables. Marital status includes the following categories: married, separated, divorced or widowed, and never married (reference category). We also control for major life events. Respondents were asked to report the occurrence of 15 major life events within the last five years, such as experiencing the death of a child/spouse or being the victim of a serious physical attack. We summed affirmative responses, with greater values indicating more stressful events. Informal social integration is measured by averaging responses to two questions about (1) the frequency of getting together with friends, neighbors, or relatives and (2) the frequency of talking on the telephone or exchanging emails with friends, neighbors, or relatives (1 = *never* to 6 = *more than once a week*) ($\alpha = 0.58$). Perceived social support comprises an average of responses to reports of the following (1 = *none of the time* to 5 = *all of the time*): have someone to confide in or talk to about yourself or your problems; have someone to take you to the doctor if you had to go; have someone to help you with your daily chores if you were sick; have someone to loan you a small amount of money ($\alpha = 0.78$).³ Finally, respondents were asked about the presence of both friends and kin members in their neighborhood: 1 = *none* to 5 = *10 or more*. The final measure of friends and kin networks averages responses to these questions.

Analytic Plan

Our analytic strategy proceeds in three stages. In the first stage, we calculate the means of everyday and major perceived discrimination for the following demographic groups: race/ethnicity, gender, income, and race/gender. We use *t* tests to compare the difference in means across groups. In the second stage, we use ordinary least squared regression models to estimate depressive symptoms, loneliness, and hostility. Everyday and major perceived discrimination are included in the first and all subsequent models. The second set of models includes the following demographic controls: age, gender, race/ethnicity, immigrant status, education, income, employment, homeownership, marital status, and living situation. The third set of models adjusts for all variables in model 2 and the following: major life events, informal social integration, perceived social support, and friend and kin networks.

Finally, in the third stage of our analyses, we examine the association between perceived discrimination and mental health for race/ethnicity, gender, income, and race/gender subgroups. We use adjusted Wald tests to compare the strength of association between perceived discrimination and mental health across social status subgroups (Brame et al. 1998).

All analyses were weighted to account for differential selection into the sample, nonresponse, and household size. With respect to age, race/ethnicity, and gender, the distribution of the weighted sample and the 2000 census estimates are comparable (Morenoff et al. 2007). Prior to data release, the CCAHS data team imputed variables missing fewer than 10 percent of observations. They used both case-by-case imputation (relying on interviewer comments when available) and a single regression-based imputation (based on demographic, economic, and psychological variables) to preserve these missing observations. The CCAHS data team did not impute household income, as 19 percent of respondents did not report their income, and we used multiple imputation to preserve these observations (IVEware via SAS, see Raghunathan et al. 2001). Our analytic sample is 3,102 respondents.⁴ All multivariate analyses were conducted using Stata's MI commands.

Sample Description

Table 1 presents weighted descriptive statistics of all variables. Respondents report an average of 3.670 on the everyday discrimination scale (range: 0 to 20) and an average of 0.672 on the major discrimination scale (range: 0 to 4). The overall sample comprises about 38 percent whites, 32 percent blacks, 26 percent Hispanics, and 4 percent other race individuals. More than half of respondents (53 percent) are women, and about two-fifths are first- or second-generation immigrants. Respondents range in age from 18 to 92 (descriptives not shown). About 28 percent of the sample is between 18 and 29 years old, 23 percent is between 30 and 39 years old, 19 percent is between 40 and 49 years old, and the rest of the respondents are 50 years old or older. In terms of SES, nearly one-fourth of respondents (23 percent) did not graduate from high school and an additional 49 percent do not have a college degree. About 64 percent are employed and 41 percent own their home. The modal respondent is married (42 percent), though 37 percent report never being married.

Table 1. Weighted Descriptive Statistics of Variables Included in Analysis (N = 3,102)

	Mean	Linearized standard error
Depressive symptoms (range: 1-4)	1.815	0.014
Loneliness (range: 1-4)	1.921	0.018
Hostility (range: 1-4)	2.524	0.017
Perceived everyday discrimination scale (range: 0-20)	3.670	0.096
Perceived major discrimination scale (range: 0-4)	0.672	0.023
Age		
18-29	0.275	0.012
30-39	0.226	0.009
40-49	0.187	0.009
50-59	0.129	0.007
60-69	0.090	0.007
70 or older	0.092	0.006
Female	0.526	0.012
Race/ethnicity		
White	0.384	0.021
Black	0.321	0.023
Hispanic	0.257	0.017
Other race	0.038	0.006
Immigrant status		
First generation	0.268	0.015
Second generation	0.137	0.009
Third generation	0.595	0.019
Education		
Less than 12 years	0.234	0.012
12 to 15 years	0.487	0.014
16 years or greater	0.279	0.018
Income		
< \$10K	0.130	0.009
\$10K to < \$30K	0.317	0.013
\$30K to < \$50K	0.227	0.011
≥ \$50K	0.327	0.015
Employed	0.644	0.012
Homeowner	0.412	0.015
Marital status		
Married	0.419	0.013
Separated	0.040	0.004
Divorced or widowed	0.175	0.008
Never married	0.367	0.013
Lives alone	0.182	0.011
Major life events (range: 0-11)	2.227	0.042
Informal social integration (range: 1-6)	4.496	0.029
Perceived social support (range: 1-5)	4.100	0.020
Friend and kin networks (range: 1-5)	2.634	0.024

RESULTS

Perceived Discrimination across Social Status Groups

Table 2 displays the weighted means of everyday and major perceived discrimination by race, gender, income, and race/gender. Consistent with prior research, blacks report more everyday and major discrimination than whites. Hispanics report more major discrimination, but not more everyday discrimination, than whites. Men report more everyday and major discrimination than women. However, the combination of race and gender is important, as black women report higher levels of discrimination than white men. There are also striking differences in everyday and major discrimination by income. Individuals with household incomes of less than \$10,000 report significantly more everyday and major discrimination than their counterparts who report greater household income.

Estimating Mental Health as a Function of Perceived Discrimination

We next estimate three indicators of mental health—depressive symptoms, loneliness, and hostility—as a function of everyday and major perceived discrimination. As displayed in Table 3, the unadjusted model shows that both everyday and major discrimination are independently associated with depressive symptoms. A one-unit increase in everyday discrimination is associated with a 0.042-point increase in depressive symptoms ($p < .001$). Similarly, a one-unit increase in major discrimination is associated with a 0.040-point increase in depressive symptoms ($p < .01$). Adjusting for demographic and socioeconomic characteristics in model 2 does little to attenuate the association between discrimination and depressive symptoms. In the final model, which includes a more extensive set of controls, the coefficient of everyday discrimination is reduced by 26 percent, though there remains a statistically significant relationship between everyday discrimination and depressive symptoms (0.031, $p < .001$). However, the association between major discrimination and depressive symptoms falls to statistical insignificance.

As in the case of depressive symptoms, higher values on the everyday discrimination scale are

Table 2. Weighted Means of Perceived Discrimination, by Social Status Groups

	Perceived everyday discrimination	Perceived major discrimination	N
Race/ethnicity			
White	3.334	0.468	983
Black	4.699***	1.013***	1,239
Hispanic	2.921	0.565*	800
Other race	3.447	0.578	80
Gender			
Male	3.934**	0.845***	1,234
Female	3.432	0.516	1,868
Income			
< \$10K	4.035	0.778	459
\$10K to < \$30K	3.531***	0.674***	1,062
\$30K to < \$50K	3.529***	0.637***	713
≥ \$50K	3.758***	0.652***	868
Race/ethnicity and gender			
White male	3.541	0.580	432
White female	3.129	0.359***	551
Black male	5.141***	1.316***	416
Black female	4.372**	0.788**	823
Hispanic male	3.236	0.761*	346
Hispanic female	2.624**	0.380**	454
Other race male	3.797	0.703	40
Other race female	2.985	0.413	40

Notes: Asterisks compare whites to blacks, Hispanics, and other race; males to females; those with income of less than \$10K to other income groups; and white males to other race/gender groups.
* $p < .05$. ** $p < .01$. *** $p < .001$.

associated with increased levels of loneliness and hostility, and this positive association persists across all three models. The final model shows that a one-unit increase in everyday discrimination is associated with a 0.030-point increase in loneliness ($p < .001$) and a 0.023-point increase in hostility ($p < .001$). Although there is no association between major discrimination and

loneliness, the association between major discrimination and hostility is statistically significant. The final model indicates that a one-unit increase in major discrimination is associated with a 0.044-point increase in hostility ($p < .01$).

One important limitation of the analyses presented is that the major discrimination scale measures *lifetime* events, and it is possible that recent experiences with major discrimination are more strongly associated with mental health than distally occurring experiences with major discrimination. Respondents were asked to report if each experience of major discrimination occurred in the past year or prior to the past year, and, in analyses not presented, we exploit the temporal variation in discrimination. Similar to the analyses presented in Table 3, that do not consider the timing of discrimination, neither recently (in the past year) nor distally (prior to the past year) occurring major discrimination is associated with depressive symptoms or loneliness in the final model. We find that recently occurring major discrimination is significantly associated with hostility ($p < .05$), and distally occurring major discrimination is marginally significantly associated with hostility ($p < .10$), but these coefficients are not statistically different from each other. Taken together, these findings may suggest that it is the occurrence and frequency of major discrimination, not the timing of major discrimination, that is associated with mental health impairments. However, it is important to note that these sensitivity analyses cannot rule out the possibility that recall bias might be occurring. Longitudinal data containing information on both current and lifetime reports of discrimination (and mental health) at multiple time points would be necessary to study the association between timing of discrimination and mental health.

Though not the central focus of this article, other coefficients are associated with mental health in ways that are consistent with expectations. Women, for example, report more depressive symptoms and less hostility than men. Blacks and Hispanics, compared to their white counterparts, report less loneliness and more hostility. Education is negatively correlated with depressive symptoms and hostility. As expected, major life events are associated with more depressive symptoms, and perceived social support is inversely related to mental health impairments.

Subgroup Variation in the Association between Perceived Discrimination and Mental Health

In Table 4, we estimate the association between discrimination and mental health separately by race, gender, and income subgroups (controlling for all covariates in model 3 of Table 3). The first panel of Table 4 presents the race-stratified results.⁵ The subgroup analyses show that the association between everyday discrimination and mental health is consistent across race/ethnicity. For whites, blacks, and Hispanics, everyday discrimination is associated with more depressive symptoms, more loneliness, and more hostility. Post hoc tests of equality (adjusted Wald tests, not presented) that compare the strength of the associations across race show that the association between everyday discrimination and hostility is stronger for Hispanics than blacks ($p < .05$). With respect to major discrimination, the subgroup analyses show that major discrimination is differentially associated with impaired mental health. There is no association between major discrimination and mental health among blacks and Hispanics, but major discrimination is associated with more loneliness (0.083, $p < .05$) and more hostility (0.074, $p < .05$) among whites. Post hoc tests of equality suggest these differences in the association between major discrimination and loneliness are at least marginally statistically significant ($p < .05$ for the difference between whites and blacks, $p < .10$ for the difference between whites and Hispanics). However, these post hoc tests of equality show no statistically significant racial variation in the association between major discrimination and hostility.

The second panel of Table 4 presents gender-stratified results. These subgroup analyses show that for both men and women, everyday discrimination is associated with worse mental health outcomes. Post hoc tests of equality show that everyday discrimination is more strongly associated with depressive symptoms ($p < .05$) and loneliness ($p < .05$) for women than for men. Women's and men's hostility is equally affected by everyday discrimination. Though women's experiences with everyday discrimination are generally more strongly linked to impaired mental health, men are more likely to suffer from major discrimination. For men but not women, major discrimination is associated with more depressive

Table 3. Ordinary Least Squares (OLS) Regression Models Estimating Mental Health Outcomes as a Function of Perceived Discrimination

	Depressive symptoms			Loneliness			Hostility		
	Model 1	Model 2	Model 3	Model 1	Model 2	Model 3	Model 1	Model 2	Model 3
Perceived everyday discrimination	0.042*** (0.003)	0.041*** (0.003)	0.031*** (0.003)	0.035*** (0.005)	0.039*** (0.005)	0.030*** (0.005)	0.024*** (0.004)	0.028*** (0.004)	0.023*** (0.004)
Perceived major discrimination	0.040** (0.014)	0.041** (0.014)	0.006 (0.014)	0.007 (0.020)	0.018 (0.020)	0.001 (0.020)	0.095*** (0.016)	0.048** (0.015)	0.044** (0.014)
Age									
18-29 (reference)									
30-39		0.038 (0.032)			0.118* (0.050)			-0.025 (0.036)	
40-49		0.068 (0.041)			0.132* (0.059)			-0.063 (0.041)	
50-59		0.050 (0.047)			0.114 (0.070)			-0.009 (0.053)	
60-69		0.074 (0.051)			0.187* (0.077)			-0.088 (0.055)	
70 or older		-0.014 (0.057)			0.182* (0.089)			-0.161** (0.059)	
Female		0.124*** (0.024)			0.039 (0.033)			-0.135*** (0.027)	
Race/ethnicity									
White (reference)									
Black		-0.042 (0.029)			-0.171*** (0.045)			-0.292*** (0.032)	
Other race		-0.003 (0.080)			0.083 (0.145)			0.142 (0.087)	
Hispanic		-0.005 (0.039)			-0.125* (0.057)			0.108** (0.040)	
Immigrant status									
First generation		-0.063 (0.038)			0.053 (0.056)			0.151*** (0.042)	
									0.137** (0.042)

(continued)

Table 3. (continued)

	Depressive symptoms			Loneliness			Hostility		
	Model 1	Model 2	Model 3	Model 1	Model 2	Model 3	Model 1	Model 2	Model 3
Second generation	-0.040 (0.041)	-0.025 (0.040)	—	-0.151** (0.056)	-0.136* (0.055)	—	0.004 (0.041)	0.009 (0.041)	—
Third generation (reference)									
Education									
Less than 12 years	0.150*** (0.040)	0.112** (0.038)	—	0.056 (0.058)	0.002 (0.057)	—	0.235*** (0.041)	0.203*** (0.040)	—
12 to 15 years	0.081** (0.031)	0.068* (0.029)	—	0.004 (0.046)	-0.010 (0.045)	—	0.136*** (0.031)	0.127*** (0.031)	—
16 years or greater (reference)									
Income									
< \$10K	0.104 (0.054)	0.084 (0.052)	—	0.143 (0.077)	0.124 (0.076)	—	0.099 (0.054)	0.087 (0.052)	—
\$10K to < \$30K	0.112** (0.036)	0.084* (0.033)	—	0.115* (0.053)	0.087 (0.051)	—	0.062 (0.041)	0.045 (0.040)	—
\$30K to < \$50K	0.020 (0.032)	0.015 (0.031)	—	0.018 (0.055)	0.014 (0.054)	—	0.017 (0.040)	0.013 (0.038)	—
≥ \$50k (reference)									
Employed	0.113*** (0.027)	-0.088** (0.027)	—	-0.089* (0.044)	-0.066 (0.043)	—	0.033 (0.033)	-0.049 (0.032)	—
Homeowner	-0.013*** (0.028)	-0.094*** (0.026)	—	-0.076 (0.041)	-0.038 (0.039)	—	0.032*** (0.032)	-0.069* (0.032)	—
Marital status									
Married	-0.065 (0.036)	-0.066 (0.035)	—	-0.135** (0.048)	-0.137** (0.047)	—	0.039 (0.039)	0.052 (0.038)	—
Separated	-0.012 (0.058)	-0.021 (0.054)	—	-0.010 (0.098)	-0.021 (0.099)	—	0.064* (0.064)	0.141* (0.063)	—
Divorced or widowed	0.046 (0.037)	0.018 (0.035)	—	0.086 (0.057)	0.075 (0.056)	—	0.041** (0.041)	0.111** (0.040)	—
Never married (reference)									
Lives alone	0.054	0.031	—	0.116*	0.080	—	0.035	-0.028	—

(continued)

Table 3. (continued)

	Depressive symptoms			Loneliness			Hostility		
	Model 1	Model 2	Model 3	Model 1	Model 2	Model 3	Model 1	Model 2	Model 3
Major life events		(0.032)	(0.031) 0.039***		(0.045)	(0.043) 0.014		(0.035)	(0.034) -0.002
Informal social integration			(0.007) -0.011			(0.010) -0.034*			(0.007) -0.026*
Perceived social support			(0.011) -0.163***			(0.016) -0.187***			(0.012) -0.106***
Friend and kin networks			(0.013) -0.006			(0.023) -0.052**			(0.017) -0.021
Constant	1.635***	1.583***	2.318***	1.788***	1.770***	2.916***	2.372***	2.204***	2.887***
R ²	0.099	0.192	0.261	0.029	0.087	0.137	0.058	0.181	0.208
N	3,102	3,102	3,102	3,102	3,102	3,102	3,102	3,102	3,102

Notes: Robust standard errors in parentheses. All analyses are weighted.

* $p < .05$. ** $p < .01$. *** $p < .001$.

Table 4. Ordinary Least Squares (OLS) Regression Models Estimating Mental Health Outcomes, by Race/Ethnicity, Gender, and Income

	Depressive symptoms	Loneliness	Hostility
Race/ethnicity			
White (n = 983)			
Perceived everyday discrimination	0.030*** (0.006)	0.024** (0.008)	0.022*** (0.006)
Perceived major discrimination	0.024 (0.026)	0.083* (0.038)	0.074* (0.030)
Black (n = 1,239)			
Perceived everyday discrimination	0.029*** (0.005)	0.023** (0.007)	0.017** (0.006)
Perceived major discrimination	0.003 (0.018)	-0.015 (0.029)	0.036 (0.019)
Hispanics (n = 800)			
Perceived everyday discrimination	0.036*** (0.007)	0.043*** (0.009)	0.033*** (0.008)
Perceived major discrimination	-0.014 (0.029)	-0.010 (0.042)	0.019 (0.032)
Gender			
Males (n = 1,234)			
Perceived everyday discrimination	0.023*** (0.004)	0.020** (0.007)	0.020*** (0.005)
Perceived major discrimination	0.046* (0.020)	0.039 (0.029)	0.067** (0.021)
Females (n = 1,868)			
Perceived everyday discrimination	0.039*** (0.004)	0.040*** (0.007)	0.028*** (0.005)
Perceived major discrimination	-0.026 (0.018)	-0.037 (0.028)	0.021 (0.020)
Income			
< \$10K (n = 459)			
Perceived everyday discrimination	0.030** (0.009)	0.027* (0.013)	0.022* (0.009)
Perceived major discrimination	0.063 (0.043)	-0.070 (0.067)	0.072 (0.041)
\$10K to < \$30K (n = 1,062)			
Perceived everyday discrimination	0.035*** (0.006)	0.027** (0.008)	0.024*** (0.006)
Perceived major discrimination	0.012 (0.028)	0.018 (0.035)	0.063* (0.030)
\$30K to < \$50K (n = 713)			
Perceived everyday discrimination	0.032*** (0.007)	0.041*** (0.010)	0.017* (0.008)
Perceived major discrimination	-0.038 (0.027)	-0.037 (0.037)	0.046 (0.032)
≥ \$50K (n = 868)			
Perceived everyday discrimination	0.028*** (0.006)	0.025** (0.009)	0.025*** (0.007)
Perceived major discrimination	0.011 (0.023)	0.037 (0.036)	0.019 (0.029)

Notes: All models include covariates from model 3 of Table 3. All analyses are weighted.

* $p < .05$. ** $p < .01$. *** $p < .001$.

Table 5. Ordinary Least Squares (OLS) Regression Models Estimating Mental Health Outcomes, by Race/Ethnicity and Gender Subgroups

	Depressive symptoms	Loneliness	Hostility
White males (<i>n</i> = 432)			
Perceived everyday discrimination	0.024** (0.008)	0.013 (0.011)	0.019* (0.009)
Perceived major discrimination	0.058 (0.037)	0.112* (0.050)	0.096* (0.046)
White females (<i>n</i> = 551)			
Everyday discrimination	0.036*** (0.008)	0.038*** (0.012)	0.025** (0.008)
Major discrimination	-0.024 (0.034)	0.046 (0.053)	0.070 (0.037)
Black males (<i>n</i> = 416)			
Everyday discrimination	0.020** (0.007)	0.019 (0.011)	0.013 (0.008)
Major discrimination	0.016 (0.026)	-0.011 (0.046)	0.040 (0.029)
Black females (<i>n</i> = 823)			
Everyday discrimination	0.036*** (0.006)	0.026** (0.010)	0.025** (0.007)
Major discrimination	-0.003 (0.025)	-0.020 (0.039)	0.033 (0.029)
Hispanic males (<i>n</i> = 346)			
Everyday discrimination	0.024** (0.009)	0.017 (0.013)	0.020 (0.010)
Major discrimination	0.025 (0.035)	0.058 (0.054)	0.047 (0.044)
Hispanic females (<i>n</i> = 454)			
Everyday discrimination	0.049*** (0.011)	0.066*** (0.014)	0.039*** (0.011)
Major discrimination	-0.040 (0.051)	-0.073 (0.063)	0.001 (0.048)

Notes: All models include covariates from model 3 of Table 3. Robust standard errors in parentheses. All analyses are weighted.

* $p < .05$. ** $p < .01$. *** $p < .001$.

symptoms (0.046, $p < .05$) and hostility (0.067, $p < .01$). Post hoc tests of equality suggest statistically significant gender differences in the association between major discrimination and depressive symptoms ($p < .05$) and between major discrimination and loneliness ($p < .10$).

Finally, the third panel of Table 4 shows little variation in the association between discrimination and mental health by income. By and large, everyday discrimination but not major discrimination is linked to less favorable mental health. There is one exception. Among those with incomes between \$10,000 and \$30,000, there is a positive association between major discrimination and hostility. Post

hoc tests of equality find no variation in the association between discrimination and mental health by income.

Variation in the Association between Perceived Discrimination and Mental Health by Race and Gender

In Table 5, we present results for the following six subgroups: white men, white women, black men, black women, Hispanic men, and Hispanic women. We find limited evidence that discrimination is negatively associated with mental health among

minority men. Among both black and Hispanic men, everyday discrimination is associated with more depressive symptoms (0.020, $p < .01$ for black men; 0.024, $p < .01$ for Hispanic men). Everyday discrimination, however, is not associated with loneliness or hostility among these subgroups, though it is associated with all three mental health outcomes for women. Additionally, we find that major discrimination is strongly linked to loneliness and hostility among white men but not white women. Post hoc tests of equality generally show that both types of perceived discrimination equally affect the mental health of these six subgroups.

DISCUSSION

We use a multiethnic, population-based sample of adults in Chicago to examine the relationship between perceived discrimination and mental health. Consistent with expectations, we find a strong, robust relationship between everyday perceived discrimination and all three indicators of mental health. Chronic stressors such as everyday discrimination can capture persistent, ongoing negative exposures, and the insidious effects of these constant indignities may produce chronic helplessness and hopelessness that may lead to depression, loneliness, or hostility (Clark et al. 1999; Williams and Mohammed 2009).

Additionally, our results point to an inconsistent association between major discrimination and mental health; major discrimination is not associated with loneliness, is not associated with depressive symptoms once we adjust for major life events and social relationships, and is robustly associated with hostility. Estimates of depressive symptoms mirror results of previous work that finds, in models controlling for both everyday and major discrimination, the relationship between major discrimination and mental health is reduced to statistical insignificance (Bennett et al. 2010; Williams et al. 1997). The fact that major discrimination is strongly associated with hostility is consistent with our expectations. Major discrimination often represents large-scale, emotionally challenging, and traumatic experiences with unfair treatment. Ethnographic research highlights how feelings of anger can emerge after being unfairly stopped and harassed by the police or being passed over for a job because of race (Feagin and Sikes 1994). These more acute experiences

may engender intense and lasting emotional reactions. Indeed, the desire for retribution that has been unmet (e.g., a police officer is not reprimanded for unfairly stopping or harassing someone due to their race) may be more salient to feelings of hostility rather than to feelings of loneliness and depression.

Taken together, these findings highlight the need for researchers to analyze discrimination comprehensively and to consider a wide array of mental health outcomes (Williams and Mohammed 2009). Both dimensions of discrimination have important, unique associations with mental health. Failing to measure both dimensions may lead to an underestimation of the influence of discriminatory stress on mental health (Williams and Mohammed 2009). It should be noted that although the magnitude of these discrimination coefficients are not large, they are similar in magnitude of more commonly studied stressors such as major life events (see Kessler et al. 1999). But the fact that discrimination remains significantly associated with mental health after the inclusion of a wide array of covariates (including demographics, social support, and stress) underscores the independent importance of discrimination as a unique social stressor.

The second goal of this article was to examine subgroup variation in the association between discrimination and mental health. By and large, the subgroup analyses and post hoc tests for equality suggest that perceived discrimination is an equal opportunity risk factor for mental health impairments among the three racial/ethnic subgroups we examined. There are a couple of exceptions, with the association between major discrimination and loneliness being stronger among whites than blacks and the association between everyday discrimination and hostility being stronger among Hispanics than blacks. The stronger associations between perceived discrimination and health for whites are not necessarily indicative of discrimination being more detrimental for the overall health of whites than blacks. As research in both stress theory and discrimination indicates, what makes discrimination harmful to vulnerable populations is the repeated and chronic experiences of discrimination over time compounded by the addition of multiple stressors that are fundamentally born out of disadvantaged social contexts (Pearlin et al. 2005; Sternthal, Slopen, and Williams 2011; Thoits 2010; Turner and Avison 2003).

Though we find little variation in the consequences of discrimination for mental health by race/ethnicity, our results provide some evidence

that the association between perceived discrimination and mental health varies by gender. We find that everyday discrimination is generally more strongly associated with mental health for women than men and that major discrimination is generally more strongly associated with mental health for men. Perhaps because experiences of acute discrimination are salient to important aspects of male identity (e.g., the ability to acquire, excel in, and maintain a job to provide for a family), these experiences might have a more profound effect on men's mental health than they do for women's mental health (Cohen 2001). Previous research shows mixed findings on differential associations between discrimination and mental health across gender, and more theoretical and empirical work (particularly in diverse samples) needs to be done to better understand the role of gender, particularly as it relates to race and class, to more fully explicate the patterns that have been found (Williams and Mohammed 2009; also see Hahm et al. 2010).

Major discrimination and mental health varies across social status subgroups, with subgroup analyses finding that major discrimination may be associated with mental health primarily for white men. However, post hoc tests of equality do not find support for this. Nevertheless, it is important to interpret these findings with discretion because they rely on small sample sizes. It is possible that major experiences with unfair treatment are particularly stressful for white men, given their privileged social status, and therefore may lead to feelings of loneliness and hostility. Unlike minority groups who have experienced historical and continued discrimination and may have learned to cope with such discrimination, it is possible that whites, particularly white men, have not developed successful coping abilities due to their relatively privileged social status. It might also be the case that lower levels of exposure to discrimination are associated with greater reactivity to it because of the novelty. Indeed, white men experience lower levels of major discrimination compared to black and Hispanic men and black women. In our sample, only 42 percent of white men reported lifetime major discrimination, compared to 74 percent of black men, 52 percent of Hispanic men, and 50 percent of black women).

In general, these findings point to the notion that perceived discrimination is an important stressor for all population groups, and a growing body of research supports the fact that experiences

of unfair treatment and daily hassles for any reason can have an impact on mental and physical health (for a review, see Williams and Mohammed 2009). Although it is clear that more work needs to be done to accurately conceptualize and assess different types of discrimination (Williams and Mohammed 2009), it may also be worthwhile to consider not only the type of stressors faced by different social status groups but to also consider that the disproportionate number of stressors faced by disadvantaged social groups might differentially affect their mental and physical health (Evans and Kim 2010; Sternthal et al. 2011; Turner and Avison 2003). For example, the poorer health of black women is not only related to experiences of discrimination but also related to social contexts of higher levels of poverty, food insecurity, life events, and neighborhood instability that combined serve to impact health (Geronimus 1992, 2001).

These findings should be interpreted with caution. For one, the cross-sectional data make it difficult to discern causal ordering. Though theoretical perspectives make it easy to imagine that perceptions of discrimination adversely affect mental health outcomes, it is possible that impaired mental health influences how a person experiences, interprets, and reacts to adverse events (Brondolo et al. 2008). A person with a negative affect may incorrectly recall the intensity and frequency of past experiences as discriminatory, even though these experiences would not be assessed similarly by a person without negative affect (Brondolo et al. 2008), or negative affect might cause others to act negatively or discriminatorily. Indeed, this is a common issue with measures of chronic stress exposure such as everyday discrimination, and this is also likely true of major discriminatory events. Though it is likely that recall accuracy may be affected by both depressed and hostile mood (particularly in a cross-sectional study), there is no research that indicates loneliness would affect recall accuracy. Given our results are generally consistent across outcomes, this lends credence to our hypotheses that the causal pathway goes from discrimination to mental health. Additionally, our findings are strengthened by the few longitudinal studies suggesting a causal effect of discrimination on mental health (e.g., Brown et al. 2000; Jackson et al. 1996; Pavalko, Mossakowski, and Hamilton 2003; Schultz et al. 2006). Similarly, research links experiences with discrimination to physical health such as hypertension, obesity, and waist circumference (Hunte

2011; Hunte and Williams 2009; Williams and Mohammed 2009), a relationship less susceptible to reverse causality. Future data collection efforts, though, should include longitudinal indicators of everyday discrimination, major discrimination, and mental health for a large, representative, and diverse sample of adults, as we cannot ascertain causal estimates with our cross-sectional data.

It should also be noted that the measure of loneliness used in the article, although rapidly becoming a standard measure of assessing this construct, focuses on the psychological manifestation of loneliness rather than objective characteristics associated with loneliness such as social isolation or lack of social integration. There is more precedent in the life events literature for studying objective characteristics of social integration as an outcome because of the impact that certain life events have on personal networks (e.g., Thoits 1982, 1984). This may help to explain why major discriminatory events are not related to loneliness. Loneliness is an indicator, but not equivalent to the aspects of social relationships that major discriminatory events would be predicted to negatively affect (e.g., loss of social networks such as loss of coworkers after being unfairly fired).

It is also possible that discrimination is measured imprecisely. Our measures of chronic and acute discrimination could be based on multiple social status characteristics. Although this measurement allows for application across demographic subgroups, we lose the comprehensiveness and nuance that other measures of discrimination might afford. For example, one reason for differences in the association between discrimination and mental health evidenced across studies may be due to the fact that discrimination has been measured a variety of ways across multiple studies (for a recent review, see Williams and Mohammed 2009). Indeed, there is no consensus on the best way to measure discrimination in studies of health (Williams and Mohammed 2009). Given our research focus on the variation in the association between discrimination and mental health across social status groups, one way to create a more comprehensive measure of discrimination is to consider including additional dimensions of discrimination, such as measures of anticipation of future occurrences of discrimination or vigilance, which may have important links to mental health in ways that differ from past experiences. The idea of vigilance as a stressor has been discussed in the public health and psychology literature (for a recent

review, see Williams and Mohammed 2009) and has also been described in ethnographic work in sociology (e.g., Feagin and Sikes 1994). Examining these relationships may also disentangle possible reasons for the association between discrimination and mental health among subgroups that experience low levels of discrimination. Since it is possible that those who experience low levels of discrimination might not anticipate the possibility of discrimination in the future (as an additional stressor) in the same ways that those who experience higher levels of discrimination would, we might see more differences in the association between perceived discrimination and health across social status subgroups. In addition, although the focus of this analysis is not solely based on race, using questions that explore the race-specific discriminatory stress such as the question, "How often do you think about your race?" developed by the CDC Measures of Racism Working Group (Jones 2002) might provide additional information and insight about reactions to more subtle and insidious forms of discrimination and institutionalized racism that might impact the relationships between experiences with discrimination and health (Jones 2001).

Despite these limitations, this study provides new insight into the consequences of discrimination for mental health in a multiethnic, population-based sample of adults. We extend prior research in the following ways: (1) by examining the independent contributions of everyday discrimination and major discrimination to mental health; (2) by considering two indicators of mental health that are not commonly considered, loneliness and hostility; and (3) by examining subgroup variation in the association between discrimination and mental health. The findings highlight the need to examine multiple indicators of both discrimination and mental health as well as to pay particular attention to both differences and similarities in the association between discrimination and mental health across race/ethnicity, gender, and socioeconomic status. Given that the association between discrimination and mental health is stronger, or similar in magnitude, for majority compared to minority racial/ethnic groups, despite the fact that reports of discrimination are higher among minority racial/ethnic groups, suggests that more research attention should be paid to the differential coping and responses to stress by race (Jackson, Knight, and Rafferty 2010; Williams and Mohammed 2009), as well as racial differences in the quantities and

magnitude of stressors faced by disadvantaged groups (Sterthal et al. 2011; Turner and Avison 2003). Finally, the findings suggest far-reaching consequences of discrimination. Future work should continue to unpack how multiple dimensions of discrimination may have implications for population health inequalities.

FUNDING

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: Both Lee's and Turney's work on this project was supported by the Robert Wood Johnson Foundation Health & Society Scholars Program. The authors thank the Robert Wood Johnson Foundation Health & Society Scholars program for its financial support. This research was also supported by Grant Nos. P50HD38986 and R01HD050467 from the National Institute of Child Health and Human Development and per the NIH Public Access Policy requires that peer-reviewed research publications generated with NIH support are made publicly available through PubMed Central. NIH is not responsible for the data collection or analyses represented in this article. The authors thank Jim House, Darrell L. Hudson, Jeff Morenoff, and the CMB working group and staff at the University of Michigan for their helpful feedback.

NOTE

1. Interpersonal discrimination is "the process by which a member, or members, of a socially defined group is, or are, treated differently (especially unfairly) because of his/her/their membership of that group" (Jary and Jary 1995:169).
2. In analyses not presented, we substitute our continuous indicators of everyday and major discrimination with dichotomous variables that indicate whether a respondent *ever* experienced discrimination. Results for the full sample are robust to this specification, but we present results using the continuous measure because we are primarily interested in the accumulation of discrimination.
3. The social support measure was derived from social support scale items included in the Americans' Changing Lives (ACL) Survey (House 2010) and is based on previous work examining the measurement and use of social support scales for health research (Cohen and Syme 1985; Cohen and Wills 1985; House et al. 1985).
4. IVEware (Imputation and Variance Estimation Software) was developed by researchers at the Survey Methodology Program, Survey Research Center, Institute for Social Research (<http://www.isr.umich.edu/src/smp/ive/>). There are some important differences between individuals who report

household income and those who do not report household income. For example, those missing data on household income are less likely to report both major discrimination and everyday discrimination. They also report fewer depressive symptoms but report similar levels of loneliness and hostility. Individuals missing data on household income are more likely to be women, less likely to be non-Hispanic black, more likely to be a first-generation immigrant, and have lower levels of education ($p < .05$). In analyses not presented, we did not impute household income but instead included a dummy variable indicating observations that were missing data on household income. Results that use this alternative strategy are substantively similar to those presented.

5. Given the small number of non-Hispanic other race individuals in the sample ($n = 80$), we do not present results for this group.

REFERENCES

- Bennett, Ian M., Jennifer F. Culhane, David A. Webb, James C. Coyne, Vijaya Hogan, Leny Mathew, and Irma T. Elo. 2010. "Perceived Discrimination and Depressive Symptoms, Smoking, and Recent Alcohol Use in Pregnancy." *Birth* 37:90-7.
- Borrell, Luisa N., C. I. Kiefe, David R. Williams, Ana V. Diez-Roux, and Penny Gordon-Larsen. 2006. "Self-reported Health, Perceived Racial Discrimination, and Skin Color in African Americans in the CARDIA Study." *Social Science & Medicine* 63:1415-27.
- Borrell, Luisa N., Ana V. Diez Roux, David R. Jacobs, Jr., Steven Shead, Sharon A. Jacksone, Sandi Shragerf, and Roger S. Blumenthal. 2010. "Perceived Racial/ethnic Discrimination, Smoking and Alcohol Consumption in the Multi-ethnic Study of Atherosclerosis (MESA)." *Preventive Medicine* 51:307-12.
- Brame, Robert, Raymond Paternoster, Paul Mazerolle, and Alex Piquero. 1998. "Testing for the Equality of Maximum Likelihood Regression Coefficients between Two Independent Equations." *Journal of Quantitative Criminology* 14:245-61.
- Brondolo, Elizabeth, Nisha Brady, Shola Thompson, Jonathan N. Tobin, Andrea Cassells, Monica Sweeney, Delano McFarlane, and Richard J. Contrada. 2008. "Perceived Racism and Negative Affect: Analyses of Trait and State Measures of Affect in a Community Sample." *Journal of Social and Clinical Psychology* 27:150-73.
- Brondolo, Elizabeth, Nisha Brady ver Halen, Melissa Pencille, Danielle Beatty, and Richard J. Contrada. 2009. "Coping with Racism: A Selective Review of the Literature and a Theoretical and Methodological Critique." *Journal of Behavioral Medicine* 32:64-88.

- Brondolo, Elizabeth, Shola Thompson, Nisha Brady, Risa Appel, Andrea Cassells, Jonathan N. Tobin, and Monica Sweeney. 2005. "The Relationship of Racism to Appraisals and Coping in a Community Sample." *Ethnicity and Disease* 15:S14-9.
- Broudy, Risa, Elizabeth Brondolo, Vonetta Coakley, Nisha Brady, Andrea Cassells, Jonathan N. Tobin, and Monica Sweeney. 2007. "Perceived Ethnic Discrimination in Relation to Daily Moods and Negative Social Interactions." *Journal of Behavioral Medicine* 30:31-43.
- Brown, George W., Mdire Ni Bhrolchain, and Tirril Harris. 1975. "Social Class and Psychiatric Disturbance among Women in an Urban Population." *Sociology* 9(2):225-54.
- Brown, George W. and Tirrel Harris. 1978. *Social Origins of Depression*. New York: Free Press.
- Brown, Tony N., David R. Williams, James S. Jackson, Harold W. Neighbors, Myriam Torres, Sherrill L. Sellers, and Kendrick T. Brown. 2000. "'Being Black and Feeling Blue': The Mental Health Consequences of Racial Discrimination." *Race and Society* 2:117-31.
- Burgos, Giovanni and Fernando I. Rivera. 2009. "The (In) Significance of Race and Discrimination among Hispanic Youth: The Case of Depressive Symptoms." *Sociological Focus* 42(2):152-71.
- Carter, Robert T. 2007. "Racism and Psychological and Emotional Injury." *The Counseling Psychologist* 35: 13-105.
- Clark, Rodney, Norman B. Anderson, Vernessa R. Clark, and David R. Williams. 1999. "Racism as a Stressor for African Americans: A Biopsychosocial Model." *American Psychologist* 54:805-16.
- Cohen, Sheldon and S. Leonard Syme. 1985. "Issues in the Study and Application of Social Support." Pp. 3-22 in *Social Support and Health*, edited by S. Cohen and S. L. Syme. San Diego, CA: Academic Press.
- Cohen, Sheldon and Thomas A. Wills. 1985. "Stress, Social Support, and the Buffering Hypothesis." *Psychological Bulletin* 98(2):310-57.
- Cohen, Theodore F. 2001. *Men and Masculinity: A Text-reader*. Belmont, CA: Wadsworth.
- Cook, Walter W. and Donald M. Medley. 1954. "Proposed Hostility and Pharisaiic-virtue Scales for the MMPI." *Journal of Applied Psychology* 38(6): 414-18.
- Coyne, James C. 1976. "Depression and the Response of Others." *Journal of Abnormal Psychology* 85:186-93.
- Dailey, Amy B., Stansislav V. Kasl, Theodore R. Holford, Tene T. Lewis, and Beth A. Jones. 2010. "Neighborhood- and Individual-level Socioeconomic Variation in Perceptions of Racial Discrimination." *Ethnicity & Health* 15:145-63.
- D'Anna, Laula H., Ninez A. Ponce, and Judith M. Siegel. 2010. "Racial and Ethnic Health Disparities: Evidence of Discrimination's Effects across the SEP Spectrum." *Ethnicity & Health* 15:121-43.
- Essed, Philomena. 1991. *Understanding Everyday Racism*. Newbury Park, CA: Sage.
- Evans, Gary W. and Pilyoung Kim. 2010. "Multiple Risk Exposure as a Potential Explanatory Mechanism for the Socioeconomic Status-health Gradient." *Annals of the New York Academy of Sciences* 1186:174-89.
- Feagin, Joe R. 1991. "The Continuing Significance of Race: Antiblack Discrimination in Public Places." *American Sociological Review* 56(1):101-16.
- Feagin, Joe R. and Karyn D. McKinney. 2003. *The Many Costs of Racism*. Lanham, MD: Rowman and Littlefield.
- Feagin, Joe R. and Melvin P. Sikes. 1994. *Living with Racism: The Black Middle-class Experience*. Boston: Beacon Press.
- Flores, Elena, Jeanne M. Tschann, Juanita M. Dimas, Elizabeth A. Bachen, Lauri A. Pasch, and Cynthia L. de Groat. 2008. "Perceived Discrimination, Perceived Stress, and Mental and Physical Health among Mexican-origin Adults." *Hispanic Journal of Behavioral Sciences* 30:401-24.
- Forman, Tyrone A. 2003. "The Social Psychological Costs of Racial Segmentation in the Workplace: A Study of African Americans' Well-being." *Journal of Health and Social Behavior* 44:332-52.
- Geronimus, Arline T. 1992. "The Weathering Hypothesis and the Health of African-American Women and Infants: Evidence and Speculations." *Ethnicity and Disease* 2(3):207-21.
- Geronimus, Arline T. 2001. "Understanding and Eliminating Racial Inequalities in Women's Health in the United States: The Role of the Weathering Conceptual Framework." *Journal of the American Medical Women's Association* 56(4):133-36.
- Gibbons, Frederick X., Paul E. Etcheverry, Michelle L. Stock, Meg Gerrard, Chih-Yuan Weng, Marc Kiviniemi, and Ross E. O'Hara. 2010. "Exploring the Link between Racial Discrimination and Substance Use: What Mediates? What Buffers?" *Journal of Personality and Social Psychology* 99: 785-801.
- Hahn, Hyeouk Chirs, Al Ozonoff, Jillian Gaumond, and Stanley Sue. 2010. "Perceived Discrimination and Health Outcome: A Gender Comparison among Asian-Americans Nationwide." *Womens Health Issues* 20(5):350-58.
- Hawley, Louise C., Mary Elizabeth Hughes, Linda J. Waite, Christopher M. Masi, Ronald A. Thisted, and John T. Cacioppo. 2008. "From Social Structural Factors to Perceptions of Relationship Quality and Loneliness: The Chicago Health, Aging, and Social Relations Study." *The Journals of Gerontology Series B: Psychological Sciences and Social Sciences* 63:S375-84.
- House, James S. 2010. *Americans' Changing Lives: Waves I, II, III, and IV, 1986, 1989, 1994, and 2002* [Computer file]. ICPSR04690-v5. Ann Arbor, MI: Inter-university Consortium for Political and

- Social Research [distributor], 2010-05-27. doi:10.3886/ICPSR04690
- House, James S., Robert L. Kahn, Jane D. McLeod, and David Williams. 1985. "Measures and Concepts of Social Support." Pp. 83-108 in *Social Support and Health*, edited by S. Cohen and S. L. Syme. San Diego, CA: Academic Press.
- Hughes, Mary Elizabeth, Linda J. Waite, Louise C. Hawley, and John T. Cacioppo. 2004. "A Short Scale for Measuring Loneliness in Large Surveys: Results from Two Population-based Studies." *Research on Aging* 26(6):655-72.
- Hunte, Haslyn E. 2011. "Association between Perceived Interpersonal Everyday Discrimination and Waist Circumference over a 9-year Period in the Midlife Development in the United States Cohort Study." *American Journal of Epidemiology* 173(11):1232-39.
- Hunte, Haslyn E. and David R. Williams. 2009. "The Association between Perceived Discrimination and Obesity in a Population-based Multiracial and Multiethnic Adult Sample." *American Journal of Public Health* 99:1285-92.
- Jackson, James S., Tony N. Brown, David R. Williams, M. Torres, S. L. Sellers, and K. Brown. 1996. "Racism and the Physical and Mental Health Status of African Americans: A Thirteen Year National Panel Study." *Ethnicity & Disease* 6:132-47.
- Jackson, James S., Katherine M. Knight, and Jane A. Rafferty. 2010. "Race and Unhealthy Behaviors: Chronic Stress, the HPA Axis, and Physical and Mental Health Disparities over the Life Course." *American Journal of Public Health* 100:933-39.
- Jary, D. and J. Jary. 1995. *Collins Dictionary of Sociology*. Glasgow: HarperCollins Publishers.
- Jones, Camara Phyllis. 2001. "Invited Commentary: 'Race,' Racism, and the Practice of Epidemiology." *American Journal of Epidemiology* 154(4):299-304.
- Jones, Camara Phyllis. 2002. Confronting Institutionalized Racism. *Phylon* 50(1/2):7-22.
- Kessler, Ronald C., Kristin D. Mickelson, and David R. Williams. 1999. "The Prevalence, Distribution, and Mental Health Correlates of Perceived Discrimination in the United States." *Journal of Health and Social Behavior* 40:208-30.
- Krieger, Nancy. 1990. "Racial and Gender Discrimination: Risk Factors for High Blood Pressure?" *Social Science & Medicine* 30:1273-81.
- McLaughlin, Katie A., Mark L. Hatzenbuehler, and Katherine M. Keyes. 2010. "Responses to Discrimination and Psychiatric Disorders among Black, Hispanic, Female, and Lesbian, Gay, and Bisexual Individuals." *American Journal of Public Health* 100:1477-84.
- Miech, Richard Allen and Micahel J. Shanahan. 2000. "Socioeconomic Status and Depression over the Life Course." *Journal of Health and Social Behavior* 41:162-76.
- Morenoff, Jeffrey D., James S. House, Ben B. Hansen, David R. Williams, George A. Kaplan, and Haslyn E. Hunte. 2007. "Understanding Social Disparities in Hypertension Prevalence, Awareness, Treatment, and Control: The Role of Neighborhood Context." *Social Science & Medicine* 65:1853-66.
- Paradies, Yin. 2006. "A Systematic Review of Empirical Research on Self-reported Racism and Health." *International Journal of Epidemiology* 35:888-901.
- Pascoe, Elizabeth A. and Laura Smart Richman. 2009. "Perceived Discrimination and Health: A Meta-analytic Review." *Psychological Bulletin* 135:531-54.
- Pavalko, Eliza K., Krysia N. Mossakowski, and Vanessa J. Hamilton. 2003. "Does Perceived Discrimination Affect Health? Longitudinal Relationships between Work Discrimination and Women's Physical and Emotional Health." *Journal of Health and Social Behavior* 44:18-33.
- Pearlin, Leonard I., Scott Schieman, Elena M. Fazio, and Stephen C. Meersman. 2005. "Stress, Health, and the Life Course: Some Conceptual Perspectives." *Journal of Health and Social Behavior* 46(2):205-19.
- Radloff, Lenore Sawyer. 1977. "The CES-D Scale: A Self-report Depression Scale for Research in the General Population." *Applied Psychological Measurement* 1:385-401.
- Raghunathan, Trivellore E., James M. Lepkowski, John Van Hoewyk, and Peter Solenberger. 2001. "A Multivariate Technique for Multiply Imputing Missing Values Using a Sequence of Regression Models." *Survey Methodology* 27(1):85-95.
- Rivera, Fernando I., Irene Lopez, Peter J. Guarnaccia, Rafael Ramirez, Glorisa Canino, and Hector Bird. 2011. "Perceived Discrimination and Antisocial Behaviors in Puerto Rican Children." *Journal of Immigrant and Minority Health* 13(3):453-61.
- Ross, Catherine E., and John Mirowsky. 1989. "Explaining the Social Patterns of Depression: Control and Problem-solving or Support and Talking." *Journal of Health and Social Behavior* 30(2):206-19.
- Sampson, Robert J., Stephen W. Raudenbush, and Felton Earls. 1997. "Neighborhoods and Violent Crime: A Multilevel Study of Collective Efficacy." *Science* 277:918-24.
- Sapolsky, Robert. 2005. "Sick of Poverty." *Scientific American* 293:92-9.
- Schwartz, Sharon and Ilan H. Meyer. 2010. "Mental Health Disparities Research: The Impact of within and between Group Analyses on Tests of Social Stress Hypotheses." *Social Science & Medicine* 70: 1111-18.
- Sternthal, Michelle J., Natalie Slopen, and David R. Williams. 2011. "Racial Disparities in Health: How Much Does Stress Really Matter?" *Du Bois Review* 8(1):95-113.

- St. Jean, Yanick and Joe R. Feagin. 1998. *Double Burden: Black Women & Everyday Racism*. Armonk: M.E. Sharpe, Inc.
- Swim, Janet K., Lauri L. Hyers, Laurie L. Cohen, and Melissa J. Ferguson. 2001. "Everyday Sexism: Evidence for Its Incidence, Nature, and Psychological Impact from Three Daily Diary Studies." *Journal of Social Issues* 57:31-53.
- Thoits, Peggy A. 1982. "Conceptual, Methodological, and Theoretical Problems in Studying Social Support as a Buffer against Life Stress." *Journal of Health and Social Behavior* 23(2):145-59.
- Thoits, Peggy A. 1984. "Explaining Distributions of Psychological Vulnerability: Lack of Social Support in the Face of Life Stress." *Social Forces* 63:453-81.
- Thoits, Peggy A. 2010. "Stress and Health: Major Findings and Policy Implications." *Journal of Health and Social Behavior* 51(1):S41-53.
- Troxel, Wendy M., Karen A. Matthews, Joyce T. Bromberger, and Kim Sutton-Tyrrell. 2003. "Chronic Stress Burden, Discrimination, and Subclinical Carotid Artery Disease in African American and Caucasian Women." *Health Psychology* 22(3):300-09.
- Turner, R. Jay and William R. Avison. 1989. "Gender and Depression: Assessing Exposure and Vulnerability in a Chronically Strained Population." *Journal of Nervous and Mental Disease* 177:433-55.
- Turner, R. Jay and William R. Avison. 2003. "Status Variations in Stress Exposure: Implications for the Interpretation of Research on Race, Socioeconomic Status, and Gender." *Journal of Health and Social Behavior* 44:488-505.
- Wethington, Elane, George W. Brown, and Ronald C. Kessler. 1995. "Interview Measurement of Stressful Life Events." Pp. 59-79 in *Measuring Stress: A Guide for Health and Social Scientists*, edited by S. Cohen, R. C. Kessler, and L. U. Gordon. New York: Oxford University Press.
- Whitbeck, Les B., Dan R. Hoyt, Barbara J. McMorris, Xiaojin Chen, and Jerry D. Stubben. 2001. "Perceived Discrimination and Early Substance Abuse among American Indian Children." *Journal of Health and Social Behavior* 42:405-24.
- Williams, David R. and Selina A. Mohammed. 2009. "Discrimination and Racial Disparities in Health: Evidence and Needed Research." *Journal of Behavioral Medicine* 32:20-47.
- Williams, David R., Harold W. Neighbors, and James S. Jackson. 2003. "Racial/ethnic Discrimination and Health: Findings from Community Studies." *American Journal of Public Health* 93:200-08.
- Williams, David R., Yan Yu, James S. Jackson, and Norman B. Anderson. 1997. "Racial Differences in Physical and Mental Health." *Journal of Health Psychology* 2:335-51.